

UNCLASSIFIED

AD NUMBER	
AD384873	
CLASSIFICATION CHANGES	
TO:	unclassified
FROM:	confidential
LIMITATION CHANGES	
TO:	Approved for public release, distribution unlimited
FROM:	Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; SEP 1967. Other requests shall be referred to Office of Naval Research, Arlington, VA 22203.
AUTHORITY	
ONR ltr 27 Jul 1991; ONR ltr 27 Jul 1991	

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED



Service through Science

Semi-Annual Progress Report No. 3

February 1, 1967 to July 31, 1967

A STUDY OF THE EXPLOSION LIMITS OF SIMPLE  
DIFLUORAMINO COMPOUNDS (U)

Contract No. Nonr-4065(00)

to

Office of Naval Research  
Washington, D. C. 20360

CLASSIFICATION

UNCLASSIFIED

CHANGED TO

AUTH. CITED IN ASTIA R/B # 69-9

DATED

3-20-69

THOMAS F. DAVIS  
(Signature & Date)

from

Kinetics and Combustion Group  
Atlantic Research Corporation  
Alexandria, Virginia

10/2 1967

Chief Investigator: G. von Elbe  
Internal Consultant: J. B. Levy  
Scientist: G. White

September, 1967

ATLANTIC  RESEARCH

UNCLASSIFIED

UNCLASSIFIED

COPY NO. 1

# ESPIONAGE CLAUSE

This document contains information affecting the national defense of the United States within the meaning of the Espionage Acts, Title 18, U.S.C., Sections 793 and 794. The transmission or the revelation of its contents in any manner to an unauthorized person is prohibited by law.

UNCLASSIFIED

TABLE OF CONTENTS

	<u>Page</u>
I. ABSTRACT . . . . .	1
II. INTRODUCTION . . . . .	2
III. STATUS OF THE PROJECT . . . . .	2
IV. PROGRESS DURING THE PRESENT PERIOD .	3
A. Analysis of Products of DP Explosions . . . . .	3
B. Branched-Chain Explosion of DP-Fluorine Mixtures . . . . .	4
V. PLANS FOR THE FUTURE . . . . .	13
VI. REFERENCES . . . . .	13

UNCLASSIFIED

UNCLASSIFIED

I. ABSTRACT

The explosion products of 1,2-DP were analyzed with the aid of a Beckman GC-2A Gas Chromatograph. For explosive conditions the volume of the major products  $N_2:C_2H_2:HCN$  was 2.5:1:1. For non-explosive decomposition conditions  $N_2$  and  $C_2H_2$  appeared in about the same quantities and HCN was slightly less than  $C_2H_2$ . The survival of thermodynamically unstable HCN and  $C_2H_2$  indicate primary reaction products, an important consideration in the formulation of the chain reaction mechanism.

$F_2$  induced explosion and inhibition of explosion by  $O_2$  is described by the reaction scheme

- (1)  $F + DP \rightarrow NF + F$   
                                   $\downarrow$  destruction
- (2)  $F + DP \rightarrow NF_2 \rightarrow$  destruction
- (3)  $F + DP + F_2 \rightarrow 2F$
- (4)  $F + DP + O_2 \rightarrow$  destruction

The postulated two modes of reaction of F and DP are reconciled with the high temperature explosion scheme of pure DP vapor.

UNCLASSIFIED

UNCLASSIFIED

## II. INTRODUCTION

This is the third semi-annual progress report on this research program. In this program we are studying the explosion limit behavior of a series of bis-difluoramino compounds. The compounds that are of interest to us are 1,2-bisdifluoraminopropane (1,2-DP); 2,2-bisdifluoraminopropane (2,2-DP); 2-methyl-1,2-bisdifluoraminopropane (IBA); 1,3-bisdifluoraminopropane (1,3-DP); 1,1-bisdifluoraminopropane (1,1-DP); and 2,3-bisdifluoraminobutane (2,3-DB).

## III. STATUS OF THE PROJECT

At the start of the present period the vapor phase explosion limit behavior of 1,2-DP, 2,2-DP, IBA, and 1,3-DP has been characterized. We have studied these limits at pressures of 1-200 torr, at temperatures between 295°C and 450°C, and vessel diameters ranging from 4.6 cm to 17 cm. In addition the effect on the limit of inert gas and the effect of the presence of one DP on the limit of another has been studied. All of the evidence presented by this data indicate that the explosion of 1,2-DP and 2,2-DP is governed by a branching chain mechanism with chain branching occurring in the gas phase and chain breaking occurring in the gas phase and at the wall. In addition, the gas-phase chain-branching and breaking reactions involve collisions of the same molecular species and are of the same order with respect to the chain carrier. The equations expressing the explosion limits are

$$\text{1,2-DP} \quad p_d^2 = \frac{1.5 \times 10^{-3} \frac{658}{T} e^{16,600/RT}}{(2e^{-900/RT} - 1)}$$

$$\text{2,2-DP} \quad p_d^2 = \frac{1.5 \times 10^{-3} \frac{568}{T} e^{14,500/RT}}{(2e^{-782/RT} - 1)}$$

UNCLASSIFIED

CONFIDENTIAL

Explosion limit data for 1,3-DP indicate a constant  $pd^2$  product. As has been explained in the previous report, this signifies that the chain-branching step is unimolecular. Furthermore, because of the absence of a low temperature limit within the pressure range studied we must eliminate the gas-phase chain breaking step and depend only on the vessel surface for chain destruction.

The equation resulting from this data for the 1,3-DP explosion limit is

$$pd^2 = 435/4.8 \times 10^{13} e^{-44,500/RT}$$

IBA explosions do not lend themselves to this type of an analysis because a self-inhibition reaction which occurs prior to explosion obscures the limit.

#### IV. PROGRESS DURING THE PRESENT PERIOD

##### A. Analysis of Products of DP Explosions

It has been previously determined that the major gaseous products of the explosion of DP are HCN,  $N_2$ ,  $C_2H_2$ , and HF. During the present period an effort was made to establish what quantitative relationship existed between these products. For this purpose we obtained a Beckman GC-2A Gas Chromatograph. The manifold containing the explosion reactor was attached to the inlet of the chromatograph. Because the pressure in the vessel after explosion is only a small fraction of an atmosphere (~50 torr) the sample was introduced through a gas-sampling valve placed before the column. The column found to be most satisfactory for this analysis was a stainless steel (thin wall), 6-ft column containing silicone 550, 30% on C-22 firebrick, 42/60 mesh. The program used was 70°C column temperature, He-flow rate 33cc/min for 2 minutes, then 86cc/min for the remainder of analysis, filament current 200 milliamp. 1,2-DP explosion products were used for these analyses. A sampling valve loop with a volume of 9.2cc was used to probe the explosion mixture.

CONFIDENTIAL

UNCLASSIFIED

Interpretation of the chromatograph curves proved difficult due to the multitude of small minor peaks, many of which shouldered the larger major peaks. Using the program described above, we obtained peaks for  $N_2$ ,  $C_2H_2$ , and HCN at 1.3, 1.45, and 1.8 minutes respectively.  $SiF_4$  could not be desorbed from the column under conditions that still gave satisfactory results for the  $N_2$ ,  $C_2H_2$ , and HCN retention times. The explosion conditions for 1,2-DP were 1 liter reactor, 18 mm of DP, and  $409^\circ C$ . The final pressure after explosion was 64 mm. The ratio  $N_2:C_2H_2:HCN$  was about 2.5:1:1. In addition, unreacted DP residing in the manifold lines outside the reactor was recovered.

When 20 mm of 1,2-DP was decomposed non-explosively in the one-liter reactor at  $384^\circ C$ ,  $N_2$  and  $C_2H_2$  appeared in about the same quantities and HCN was slightly less than  $C_2H_2$ . We conclude that there is no fundamental difference in the over-all chemistry of the explosive and non-explosive reaction. The significant fact is the survival of  $C_2H_2$  and HCN as major reaction products. Since both compounds are thermodynamically unstable at the temperatures and pressures of these experiments, they evidently represent primary decomposition products of the DP-molecule. This is an important consideration in the formulation of the chain reaction mechanism, as has already been noted in earlier reports. The larger yield of  $N_2$  in the explosion experiments does not seem to be significant. It may be expected that less  $C_2H_2$  and HCN survive in the explosion transient than in the non-explosive reaction, so that more of the stable products  $N_2$ , carbon and  $H_2$  are formed. We did not attempt to determine yields of the two latter products. No trace of fluorocarbon was found. It appears that fluorine reacts to form HF which in turn reacts with the glass reaction vessel to form  $SiF_4$ .

#### B. Branched-Chain Explosion of DP-Fluorine Mixtures

It was found that mixtures of DP and fluorine explode spontaneously at low temperatures when the fluorine concentration exceeds a critical limit. This discovery resulted from attempts to prepare mixtures of DP and fluorine at room temperature. For this purpose fluorine was gradually added in small increments to 1,2-DP vapor. The reaction vessel was a 500cc Pyrex bulb and the Pyrex inlet tube was provided with two stopcocks in series, so as to

UNCLASSIFIED



UNCLASSIFIED

leave a space of 3cc between the stopcocks. Fluorine was admitted to this space at a pressure exceeding the 1,2-DP pressure in the vessel; in this way it was possible to admit a small amount of fluorine to the reactor by closing the upper stopcock and opening the lower stopcock. Table 1 summarizes the data and observations.

TABLE 1

Fluorine-Induced Explosion of 1,2-DP Vapor at  
Room Temperature (Experiments performed by  
Mr. Wesley Copeland.)

<u>Experiment</u>	<u>Pressure, mmHg</u>		<u>Comments</u>
	<u>1,2-DP</u>	<u>1,2-DP + F<sub>2</sub></u>	
1	20mm	20.9	F <sub>2</sub> added in increments of app. 0.25mm, no visible reaction after standing 2 hrs.
2	22	22.9	F <sub>2</sub> added in increments of 0.45mm, no reaction.
		23.5	The addition of 0.6mm more of F <sub>2</sub> produced an explosion. Final pressure 61mm.
3	18.5	19.7	1.2mm of F <sub>2</sub> put in, in one shot. No visible reaction.
		20.15	0.45mm more F <sub>2</sub> added - explosion; final pressure, 51mm +
		51 + .66	0.66mm more F <sub>2</sub> added - another smaller explosion.
4	19.5	21.0	1.5mm of F <sub>2</sub> added in one shot - small flame in bore of stopcock but no explosion. Final pressure 21.0mm
		21.3	0.3mm more of F <sub>2</sub> put in, in 0.15mm shots. No visible reaction.

By infrared analysis it was shown that the explosion products contain HCN and C<sub>2</sub>H<sub>2</sub>. SiF<sub>4</sub> and CF<sub>4</sub> were also identified.

UNCLASSIFIED

CONFIDENTIAL

Numerous additional data were obtained using the following apparatus and experimental procedure (Fig. 1). A vacuum line made up of teflon tubing, fittings, and valves was built to handle the fluorine experiments. Inlets were provided for the DP, the additive gases as well as fluorine. A glass capillary and bulb mercury manometer measured the pressures in the line. The bulb was coated with a layer of Kel-F wax and the mercury with Kel-F oil. A glass reactor was fabricated with separate inlets for DP and for the additive gases in order to prevent mixing of the DP and the additives in the stopcock bore. Pressures in the reactor volume were measured by a U-tube mercury manometer. After the system was completely evacuated a measured amount of DP was led into the reactor with stopcocks (1) and (3) open and (2) closed. Additive gases fill the line with (2) and (3) closed and (1) open to a pressure determined by the ratio of volumes defined by the three stopcocks and the reactor volume according to the formula

$$\% \text{ additive} = \frac{v}{V} \left( \frac{\frac{p}{P} - 1}{\frac{v}{V} \frac{p}{P} + 1} \right) \times 100$$

Where  $p$  and  $v$  denote the pressure and volume between stopcocks 1, 2, and 3, and  $P$  and  $V$  denote the pressure volume of the reactor. Table 2 contains the data obtained in these experiments.




Fig. 1. Scheme of Apparatus for Admitting  $F_2$  to DP Vapor

CONFIDENTIAL

# CONFIDENTIAL

TABLE 2

Critical Pressure of Fluorine at Fluorine-Induced DP-Explosion Limit. Effect of DP-Pressure, Temperature, Vessel Dimension and Additives

	DP Pressure torr	Temp. °C	Volume cc	Pressure O <sub>2</sub> torr	Pressure N <sub>2</sub> torr	Pressure A torr	Pressure F <sub>2</sub> at limit torr
1,2 DP	8	25	200	-	-	-	2.3, 2.5→2.8, 2.3
	20	25	200	-	-	-	2.0
	8	55	200	-	-	-	2.5
	8	1	200	-	-	-	3.5
	8	8	200	-	-	-	3.0
	8	100	200	-	-	-	1.24
	8	25	1000	-	-	-	1.40
	8	25	200	2.2	-	-	6.6→6.9, 4.8
	8	25	200	2.2	-	9.8	7.0→7.3
	8	25	200	-	2.3	-	2.3
	8	25	200	1.1	-	-	3.5
	8	25	200	3.6	-	-	8.0
2,2 DP	8	25	200	-	-	-	2.8→3.4
IBA	8	25	200	-	-	-	< 0.2
	8	25	1000	-	-	-	0.58
	8	25	200	-	2.3	-	0.4
	8	25	200	5.0, 7.4, 2.3	-	-	0.4
1,3 DP	8	25	200	-	-	-	0.4
	8	25	200	2.2	-	-	3.0→3.3, 4.6
	8	25	200	2.2	-	9.2	4.1
	8	25	200	2.2	2.1	-	3.9
	8	25	200	1.1	-	-	1.0

CONFIDENTIAL

From the data in Tables 1 and 2 we draw the following general conclusions:


(1) Within the range of experimental scatter the critical fluorine pressure is independent of the pressure of DP vapor, the vessel dimension and inert-gas addition.

(2) There is a small decrease of the critical fluorine pressure over the temperature range from 1° to 100°C.

(3) Addition of oxygen causes a substantial increase of the critical fluorine pressure; this increase is unaffected by simultaneous addition of inert gas.

The experimental facts rule out *prima facie* any thermal interpretation of the explosion phenomenon, which is thus identified as a branched-chain explosion. In performing these experiments we have unfortunately not been aware of an obvious inference from fact 3; namely, since oxygen is an inhibitor in these explosions, and since elementary fluorine may react slightly with the glass wall of the apparatus to liberate oxygen, there exists the probability that all these data are slightly falsified by traces of oxygen. Experiments are planned to test this point; we feel, however, that despite this uncertainty the present data are sufficiently definitive to permit valid conclusions concerning the kinetics and chemical mechanism of the fluorine-induced explosive reaction.

At the explosion limit the rate of chain breaking equals the rate of chain branching. Fact 1 signifies that both chain breaking and chain branching occur in the gas phase; that both rates are of the same order with respect to the DP-concentration; and that chain branching involves collisions with  $F_2$ -molecules. Thus, if X denotes a chain carrier, the reaction scheme might be of the general type



so that the explosion condition becomes

$$k_a [DP][X] = k_b [DP][F_2][X]$$




CONFIDENTIAL

CONFIDENTIAL

or

$$[F_2] = k_a/k_b \quad (1)$$

Alternatively, the scheme might be



which yields again  $[F_2] = k_a/k_b$ . The difference is that in the former case reaction (a) is bimolecular and reaction (b) is trimolecular; in the latter case the reactions are unimolecular and bimolecular, respectively.

Typical or "ballpark" values of the pre-exponential factors of rate coefficients may be quoted as

$10^{12} \text{ sec}^{-1}$	for unimolecular reactions
$10^{-11} \text{ cm}^3 \text{ sec}^{-1}$	for bimolecular reactions
$10^{-34} \text{ cm}^6 \text{ sec}^{-1}$	for trimolecular reactions

Thus, for both cases, roughly,

$$[F_2] \approx 10^{23} e^{-(E_a - E_b)/RT} \quad (2)$$

and since for 1,2-DP at room temperature according to Tables 1 and 2,

$$[F_2] \approx 2 \text{ mmHg} \approx .7 \times 10^{17} \text{ molecules/cm}^3,$$


one calculates the activation energy  $E_a - E_b$  to be approximately 8,500 cal/mole. On this basis,  $[F_2]$  should increase over the temperature range from 1°C to 100°C by a factor of approximately 80. However, according to fact 2  $[F_2]$  does not increase at all but decreases instead over this temperature range.

We may introduce a "steric" factor to reduce the value of the pre-exponential factor of reaction (a); but in order to accommodate the theory to fact 2 the steric factor would have to be absurdly small, so as to be totally implausible on the basis of collision concepts.

CONFIDENTIAL

$$(1) \quad \text{NF}_2 + \text{DP} \rightarrow \text{HCN} + \text{N}_2 + \text{C}_2\text{H}_2 + 3\text{HF} + 3\text{F}$$
$$(2) \quad \text{NF}_3 + \text{DP} \rightarrow \text{HCN} + \text{N}_2 + \text{C}_2\text{H} + 4\text{HF} + \text{F}_2$$


We propose that the key to the situation is to be found in the molecular rearrangements that occur upon attack of a DP-molecule by the F-atom. This attack results in bonding of F with one of the H-atoms of the DP-molecule, with a net exothermicity of approximately 55 kcal/mole. The C-F bond thus generated will then preferentially engage one of the DP-groups, forming a C-N bond of bond energy  $\approx 55$  kcal/mole and breaking one of the DP N-F bonds ( $\approx 70$  kcal/mole) by partial utilization of the 55 kcal/mole available from the F-atom attack. Alternatively, but much less frequently, the C-F bond engages the adjacent C-atom, forming a double bond C=C and leaving an  $\text{NF}_2$ -group. Thus,



10

UNCLASSIFIED

explosion is not  $\text{NF}_2$  but  $\text{NF}$ . Thus, the high-temperature explosion scheme becomes




and the explosion boundary condition becomes (with  $k_5 \gg k_6$ )

$$\frac{2}{k_2} \frac{k_1}{k_5} - 1 = \frac{k_3}{k_2 [\text{DP}]} \quad (3)$$

which is of the same form as the former equation (1) in Semi-Annual Report No. 1, and permits substantially the same analysis of the data. We have merely removed the logical difficulty of attributing nearly identical activation energies to the collision processes (1) and (2) and have introduced instead the concept of two different modes of break-up of the energized DP-radical, one mode being much more frequent than the other.

With respect to the  $\text{F}_2$ -induced low-temperature explosion, we admit the additional ternary reaction



which implies that the transient (DP-radical +  $\text{F}_2$ ) collision complex is sufficiently energized to permit the exchange  $\text{NF} + \text{F}_2 = \text{NF}_2 + \text{F}$ . Furthermore, the kinetic analysis of the high-temperature explosion limit leads to the conclusion that at the low temperatures of the fluorine experiments the radicals  $\text{NF}$  and  $\text{NF}_2$  do not significantly react with DP. From literature data

UNCLASSIFIED

UNCLASSIFIED


(ref. 1) it can also be shown that under the conditions of these experiments these radicals do not significantly react in collisions with  $F_2$ . Thus, the fluorine-induced explosions are described by reactions (3), (4), (5), (6) and (7), yielding the explosion boundary condition

$$[F_2] = \frac{k_4}{k_7} \frac{k_6}{k_5} \quad (4)$$

On the basis of the arguments presented in connection with equations (1) and (2), we conclude that reactions (4) and (7) have very low activation energies and that the ratio  $k_5/k_6$  is of the order of  $10^6$ .

Returning to equation (3), if the difference of activation energies between reactions (1) and (2) were of the order of the dissociation energy of  $F_2$ , i.e., 35.6 kcal/mole,  $k_5/k_6$  would be of the order of  $10^{11}$ . The temperature is here about 400°C as compared to room temperature in the fluorine experiments. This temperature difference may be a factor in the disparity of the values of the ratio  $k_5/k_6$ ; it is, however, reasonable to assume that the difference of activation energies between reaction (1) and (2) is considerably less than the  $F_2$ -dissociation energy, owing to intramolecular energy available in the break-up of the collision complex (NF + DP); and hence the disparity of the values  $k_5/k_6$  is considerably diminished.

The inhibition of the fluorine-induced explosion by oxygen is explained quite simply by the additional ternary reaction



whose detailed chemistry may be imagined in many ways. The explosion condition becomes

$$[F_2] = \frac{k_4}{k_7} \frac{k_6}{k_5} + \frac{k_8}{k_7} [O_2] \quad (5)$$

which is of a form that appears to be consistent with the data.

UNCLASSIFIED



UNCLASSIFIED

V. PLANS FOR THE FUTURE

Further chemical and kinetic studies are planned with a view of refining the concept of free-radical reactions and intramolecular rearrangements of difluoramino compounds, to yield active and inert molecular fragments. Attention will be given to effects of molecular structure, including structures comprising more than three carbon atoms in the molecule, and to coordination of our data and concepts with work performed elsewhere, particularly at S.R.I. Additional experiments on fluorine-induced explosions are included in this program. A special effort will be made to eliminate any fortuitous inhibition of the reaction by oxygen contamination.

We expect that during the current period the work on difluoramino propanes will be substantially completed and significant data on the butane members of the series will be obtained.

VI. REFERENCES

1. J. B. Levy and B. K. Wesley Copeland, J. Phys. Chem. 69, 3700 (1965).

UNCLASSIFIED

~~UNCLASSIFIED~~

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1 ORIGINATING ACTIVITY (Corporate author) Atlantic Research Corporation Shirley Highway and Edsall Road Alexandria, Virginia		7a REPORT SECURITY CLASSIFICATION CONFIDENTIAL
		2b GROUP
3 REPORT TITLE A STUDY OF THE EXPLOSION LIMITS OF SIMPLE DIFLUORAMINO COMPOUNDS (U)		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Semi-Annual Progress Report No. 3, February 1, 1967 to July 31, 1967		
5 AUTHOR(S) (Last name, first name, initial) von Elbe, Guenther Levy, J. B. White, G.		
6. REPORT DATE September, 1967	7a. TOTAL NO OF PAGES 13	7b. NO. OF REFS 1
8a CONTRACT OR GRANT NO. Nonr-4065 (00)	9a. ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO.		
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10 AVAILABILITY/LIMITATION NOTICES		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Office of Naval Research Washington, D. C. 20360	
13 ABSTRACT <p>The explosion products of 1,2-DP were analyzed with the aid of a Beckman GC-2A Gas Chromatograph. For explosive conditions the volume of the major products <math>N_2:C_2H_2:HCN</math> was 2.5:1:1. For non-explosive decomposition conditions <math>N_2</math> and <math>C_2H_2</math> appeared in about the same quantities and HCN was slightly less than <math>C_2H_2</math>. The survival of thermodynamically unstable HCN and <math>C_2H_2</math> indicate primary reaction products, an important consideration in the formulation of the chain reaction mechanism.</p> <p><math>F_2</math> induced explosion and inhibition of explosion by <math>O_2</math> is described by a chemical reaction scheme. The postulated reaction scheme of F and DP is reconciled with the high temperature explosion scheme of pure DP vapor.</p>		

DD FORM 1473  
1 JAN 64

~~UNCLASSIFIED~~

Security Classification

~~UNCLASSIFIED~~

Security Classification

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	bisdifluoraminopropane chromatograph explosion chain reaction fluorine inhibition oxygen						

#### INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION.** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER.** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S).** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY.** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT.** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS) (S) (C) or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS.** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

~~UNCLASSIFIED~~

Security Classification